SPECIMEN LABEL Avid® 0.15EC



Miticide/Insecticide

For control of leafminers and mites and suppression of aphids, whiteflies, and thrips on ornamental plants

Recommended for Agricultural/Commercial Use

Active Ingredients:

Abamectin (CAS No. 65195-56-4 and 65195-55-3) 2.0*%

*1 gal. contains 0.15 lb. abamectin

KEEP OUT OF REACH OF CHILDREN. WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See additional precautionary statements and directions for use inside booklet.

EPA Reg. No. 100-896 EPA Est. 39578-TX-001 SCP 896A-L6A 0904 1 gallon U.S. Standard Measure



FIRST AID		
If swallowed	Call poison control center or doctor immediately for treatment advice. Do not give any liquid to the person. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.	
If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.	
If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.	
If inhaled	 Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. 	

NOTE TO PHYSICIAN

Early signs of intoxication include dilation of pupils, muscular incoordination, and muscular tremors. Toxicity following accidental ingestion of Avid can be minimized by early administration of chemical adsorbents (e.g., activated charcoal).

If toxicity from exposure has progressed to cause severe vomiting, the extent of resultant fluid and electrolyte imbalance should be gauged. Appropriate supportive parental fluid replacement therapy should be given, along with other required supportive measures (such as maintenance of blood pressure levels and proper respiratory functionality) as indicated by clinical signs, symptoms, and measurements.

In severe cases, observations should continue for at least several days until clinical condition is stable and normal. Since abamectin is believed to enhance GABA activity in animals, it is probably wise to avoid drugs that enhance GABA activity (barbiturates, benzodiazepines, valproic acid) in patients with potentially toxic

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

HOT LINE NUMBER

For 24-Hour Medical Emergency Assistance (Human or Animal) Or Chemical Emergency Assistance (Spill, Leak, Fire or Accident)

1-800-888-8372

SPECIMEN LABEL Avid® 0.15EC 2

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals WARNING

Causes substantial, but temporary eye injury. Do not get in eyes or on clothing. Prolonged or frequently repeated exposure may cause allergic skin reactions in some individuals. Harmful if inhaled or absorbed through the skin. Do not breathe spray mist. Avoid contact with skin. May be fatal if swallowed.

Personal Protective Equipment

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category B on an EPA chemical-resistance category selection chart

Applicators and other handlers must wear:

- · Long-sleeved shirt and long pants
- Chemical-resistant gloves—EPA chemical-resistance category B (e.g., barrier laminate or butyl rubber ≥14 mils).
- Shoes plus socks
- Protective eyewear

For shadehouse and greenhouse uses, applicators and other handlers must wear a dust/mist filtering NIOSH approved respirator with any R, P, or HE filter.

Wear chemical-resistant headgear for overhead exposure. Wear chemical-resistant apron when cleaning equipment, mixing, or loading. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

This pesticide is toxic to fish and wildlife. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Physical or Chemical Hazards

Do not use or store near heat or open flame.

ATTENTION

This product contains a chemical (N-methyl pyrrolidone) known to the state of California to cause birth defects or other reproductive harm

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, Inc. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and Buyer and User assume the risk of any such use. SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall SYNGENTA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

DIRECTIONS FOR USE

It is a violation of Federal (U.S.A) law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves—EPA chemical-resistance category B (e.g., barrier laminate or butyl rubber ≥14 mils).
- Shoes plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure

FAILURE TO FOLLOW DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR INSECT CONTROL, AND/OR ILLEGAL RESIDUES.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not reenter treated areas until sprays have dried.

SPECIMEN LABEL Avid® 0.15EC 3

GENERAL INFORMATION

Avid 0.15EC is an emulsifiable concentrate containing 0.15 lb. abamectin/gal. which when mixed with water according to the directions for use will control leafminers and mites and suppress whiteflies, thrips, and aphids on ornamentals.

Mix with sufficient water and apply as a foliar spray to obtain uniform coverage. For mites, apply when mites first appear and repeat as necessary to maintain control. For leafminers, apply as needed and repeat at 7-day intervals or as necessary to maintain control.

For suppression of aphids, thrips, and whiteflies apply when young, immature stages of these pests are first observed and repeat every 7 days for 2 or 3 weeks. After which time, rotate to other products that have different modes of action than Avid for at least 2-3 weeks. Refer to the **Resistance Management** section for additional comments on rotation. Aphids, thrips, and whiteflies are killed by direct contact with the Avid spray.

Residual control of pests may be enhanced with the addition of a horticultural spray oil at 0.5 to 1.0% of the spray volume on field-grown woody ornamentals, landscape plants, and Christmas trees. Repeat application as necessary, but no sooner than 7 days to maintain control. Some plants are sensitive to oils and so without prior experience the user should spray a small number of plants and observe plants for 2 weeks before spraying the remaining plants. Excessive cold or warm temperatures may increase the chance of plant damage following application with oils. Carefully read and follow directions on the oil label and do not exceed maximum rates listed on either label.

Resistance Management

Treatment may not be effective against these pests if Avid tolerant strains develop. Therefore, when applying Avid to plants that are hosts of labeled pests with multiple generations per crop or year, resistance management strategies must be used. Such strategies may include, but are not limited to, rotation of products with different modes of action, avoid treatment of successive generations with the same product, use of labeled rates at specified spray intervals, non-chemical alternatives such as beneficial arthropods, rotation of susceptible to non-susceptible plants, and various cultural practices. For additional information on the implementation of these or other resistance management strategies, consult with your State Cooperative Extension Service.

To manage susceptibility in *Liriomyza* leafminer species, apply a maximum of three applications of Avid and then rotate to Citation® for a maximum of three applications, before rotating back to Avid or using another product. The rotation between these two products with different modes of action should be based on the generation time of *Liriomyza* specie to avoid applying Avid to successive generations.

Restrictions

Do not uses Avid for suppression of aphids, whiteflies, and thrips on roses, chrysanthemums, and gerbera. These ornamentals are primary hosts of mites and *Liriomyza* leafminers for which Avid applications should be targeted. Additional applications of Avid to suppress aphids, thrips, and whiteflies on these plants will increase the selection pressure on mites and *Liriomyza* leafminers which may result in greater tolerance to Avid among these pests.

Do not apply this product through any type of irrigation system. Not for aerial application.

Do not use in citrus nurseries.

Avid has been evaluated for phytotoxicity on a wide range of ornamental plants. However, since all combinations or sequences of pesticide sprays including surfactants and adjuvants have not been tested, it is recommended that a small area be sprayed first to make certain that no phytotoxicity occurs. Phytotoxicity has been observed following the use of Avid on certain species of ferns (e.g., *Adiantum* spp.) and Shasta Daisy (*L. eucanthemum* spp.). It is therefore recommended that Avid not be used on ferns or Shasta Daisy.

CROP USE DIRECTIONS

Shadehouse, Greenhouse, Field-Grown Ornamentals, Foliage Plants, Christmas Trees, and Other Woody Ornamentals

Pests	fl. oz./100 gal.	Comments
Mites: ¹ European Red Mite Twospotted Spider Mite Camine Spider Mite Southern Red Mite Spruce Spider Mite Tarsoneimid Mites ³ Cyclamen and Broad Mites Eriophyid Mites: Rust and Bud Mites	4	³ For tarsonemid mites, repeat applications to newly developing tissue may be necessary to maintain control.
<i>Liriomyza</i> Leafminers ²	8	Repeat at 7-day intervals or as necessary to maintain control.
Boxwood Leafminer	8	For control of mining larvae, make the application when adults are beginning to lay eggs in the new foliage.
Aphids, Thrips, and Whiteflies	8	For suppression of pest populations, young immatures must be contacted by the spray.

General Information Per Application: Do not apply less than 8 fl. oz. or more than 16 fl. oz./A. Use sufficient water to obtain uniform plant coverage. Refer to the Resistance Management section and the Restrictions section for additional information.

STORAGE AND DISPOSAL

Storage

Store in a tightly closed container in a cool, dry place.

Prohibitions

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal

Triple rinse (or equivalent) and dispose of in an incinerator or landfill approved for pesticide containers.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

 $\mathsf{Avid}^{\circledcirc}, \; \mathsf{Citation}^{\circledcirc} \; \mathsf{and} \; \mathsf{the} \; \mathsf{Syngenta} \; \mathsf{logo} \; \mathsf{are} \; \mathsf{trademarks} \; \mathsf{of} \; \mathsf{a} \; \mathsf{Syngenta} \; \mathsf{Group} \; \mathsf{Company}$

U.S. Patent No. 4,310,519

©2004 Syngenta

For non-emergency (e.g. current product information) call Syngenta Crop Protection at 1-800-334-9481.

Syngenta Crop Protection, Inc. Greensboro, North Carolina 27409

www.syngenta-us.com

SCP 896A-L6A 0904

¹Apply, for example, in 200-400 gals. of water/A. In volumes of water below 200 gals., use a minimum of 8 fl. oz./A. If more than 400 gals. of water/A are required for good plant coverage, apply a maximum rate of 16 fl. oz./A. For example, if 650 gals. of water are required, use 2.5 fl. oz./100 gals.

²Apply, for example, in 100-200 gals. of water/A. In water volumes below 100 gals., use a minimum of 8 fl. oz./A. If more than 200 gals. of water per acre are required for good plant coverage, apply the maximum rate of 16 fl. oz./A. For example, if 400 gals. of water are required, use 4 fl. oz./100 gals.